

# NAG Library Function Document

## nag\_numdiff\_1d\_real\_absci (d04bbc)

### 1 Purpose

nag\_numdiff\_1d\_real\_absci (d04bbc) generates abscissae about a target abscissa  $x_0$  for use in a subsequent call to nag\_numdiff\_1d\_real\_eval (d04bac).

### 2 Specification

```
#include <nag.h>
#include <nagd04.h>

void nag_numdiff_1d_real_absci (double x_0, double hbase, double xval[])
```

### 3 Description

nag\_numdiff\_1d\_real\_absci (d04bbc) may be used to generate the necessary abscissae about a target abscissa  $x_0$  for the calculation of derivatives using nag\_numdiff\_1d\_real\_eval (d04bac).

For a given  $x_0$  and  $h$ , the abscissae correspond to the set  $\{x_0, x_0 \pm (2j - 1)h\}$ , for  $j = 1, 2, \dots, 10$ . These 21 points will be returned in ascending order in **xval**. In particular, **xval**[10] will be equal to  $x_0$ .

### 4 References

Lyness J N and Moler C B (1969) Generalised Romberg methods for integrals of derivatives *Numer. Math.* **14** 1–14

### 5 Arguments

- |    |  |               |
|----|--|---------------|
| 1: | <b>x_0</b> – double<br><i>On entry:</i> the abscissa $x_0$ at which derivatives are required.  | <i>Input</i>  |
| 2: | <b>hbase</b> – double<br><i>On entry:</i> the chosen step size $h$ . If $h < 10\epsilon$ , where $\epsilon = \text{nag\_machine\_precision}$ , then the default $h = \epsilon^{(1/4)}$ will be used. | <i>Input</i>  |
| 3: | <b>xval</b> [21] – double<br><i>On exit:</i> the abscissae for passing to nag_numdiff_1d_real_eval (d04bac).   | <i>Output</i> |

### 6 Error Indicators and Warnings

None.

### 7 Accuracy

Not applicable.

### 8 Parallelism and Performance

Not applicable.

## 9 Further Comments

The results computed by `nag_numdiff_1d_real_eval` (d04bac) depend very critically on the choice of the user-supplied step length  $h$ . The overall accuracy is diminished as  $h$  becomes small (because of the effect of round-off error) and as  $h$  becomes large (because the discretization error also becomes large). If the process of calculating derivatives is repeated four or five times with different values of  $h$  one can find a reasonably good value. A process in which the value of  $h$  is successively halved (or doubled) is usually quite effective. Experience has shown that in cases in which the Taylor series for the objective function about  $x_0$  has a finite radius of convergence  $R$ , the choices of  $h > R/19$  are not likely to lead to good results. In this case some function values lie outside the circle of convergence.

## 10 Example

See Section 10 in `nag_numdiff_1d_real_eval` (d04bac).

---