

# NAG Library Chapter Introduction

## F02 – Eigenvalues and Eigenvectors

### Contents

<b>1</b>	<b>Scope of the Chapter</b> .....	2
<b>2</b>	<b>Background to the Problems</b> .....	2
2.1	Standard Eigenvalue Problems .....	2
2.1.1	Standard symmetric eigenvalue problems .....	2
2.1.2	Standard nonsymmetric eigenvalue problems .....	3
2.2	The Singular Value Decomposition .....	4
2.3	Generalized Eigenvalue Problems.....	5
2.3.1	Generalized symmetric-definite eigenvalue problems.....	5
2.3.2	Generalized nonsymmetric eigenvalue problems .....	6
2.4	Quadratic eigenvalue problems .....	7
<b>3</b>	<b>Recommendations on Choice and Use of Available Routines</b> .....	8
3.1	Black Box Routines and General Purpose Routines .....	8
3.2	Computing Selected Eigenvalues and Eigenvectors .....	8
3.3	Storage Schemes for Symmetric Matrices.....	8
3.4	Balancing for Nonsymmetric Eigenproblems.....	9
3.5	Non-uniqueness of Eigenvectors and Singular Vectors .....	9
<b>4</b>	<b>Decision Trees</b> .....	9
4.1	Black Box Routines.....	9
4.2	General Purpose Routines (Eigenvalues and Eigenvectors).....	11
4.3	General Purpose Routines (Singular Value Decomposition) .....	12
<b>5</b>	<b>Functionality Index</b> .....	12
<b>6</b>	<b>Auxiliary Routines Associated with Library Routine Parameters</b> .....	12
<b>7</b>	<b>Routines Withdrawn or Scheduled for Withdrawal</b> .....	13
<b>8</b>	<b>References</b> .....	14

## 1 Scope of the Chapter

This chapter provides routines for various types of matrix eigenvalue problem:

- standard eigenvalue problems (finding eigenvalues and eigenvectors of a square matrix  $A$ );
- singular value problems (finding singular values and singular vectors of a rectangular matrix  $A$ );
- generalized eigenvalue problems (finding eigenvalues and eigenvectors of a matrix pencil  $A - \lambda B$ );
- quadratic eigenvalue problems (finding eigenvalues and eigenvectors of the quadratic  $\lambda^2 A + \lambda B + C$ ).

Routines are provided for both real and complex data.

The majority of routines for these problems can be found in Chapter F08 which contains software derived from LAPACK (see Anderson *et al.* (1999)). However, you should read the introduction to this chapter before turning to Chapter F08, especially if you are a new user. Chapter F12 contains routines for large sparse eigenvalue problems, although one such routine is also available in this chapter.

Chapters F02 and F08 contain **Black Box** (or **Driver**) routines that enable many problems to be solved by a call to a single routine, and the decision trees in Section 4 direct you to the most appropriate routines in Chapters F02 and F08. The Chapter F02 routines call routines in Chapters F07 and F08 wherever possible to perform the computations, and there are pointers in Section 4 to the relevant decision trees in Chapter F08.

## 2 Background to the Problems

Here we describe the different types of problem which can be tackled by the routines in this chapter, and give a brief outline of the methods used to solve them. If you have one specific type of problem to solve, you need only read the relevant sub-section and then turn to Section 3. Consult a standard textbook for a more thorough discussion, for example Golub and Van Loan (1996) or Parlett (1998).

In each sub-section, we first describe the problem in terms of real matrices. The changes needed to adapt the discussion to complex matrices are usually simple and obvious: a matrix transpose such as  $Q^T$  must be replaced by its conjugate transpose  $Q^H$ ; symmetric matrices must be replaced by Hermitian matrices, and orthogonal matrices by unitary matrices. Any additional changes are noted at the end of the sub-section.

### 2.1 Standard Eigenvalue Problems

Let  $A$  be a square matrix of order  $n$ . The *standard eigenvalue problem* is to find eigenvalues,  $\lambda$ , and corresponding eigenvectors,  $x \neq 0$ , such that

$$Ax = \lambda x. \quad (1)$$

(The phrase ‘eigenvalue problem’ is sometimes abbreviated to *eigenproblem*.)

#### 2.1.1 Standard symmetric eigenvalue problems

If  $A$  is real symmetric, the eigenvalue problem has many desirable features, and it is advisable to take advantage of symmetry whenever possible.

The eigenvalues  $\lambda$  are all real, and the eigenvectors can be chosen to be mutually orthogonal. That is, we can write

$$Az_i = \lambda_i z_i \quad \text{for } i = 1, 2, \dots, n$$

or equivalently:

$$AZ = ZA \quad (2)$$

where  $A$  is a real diagonal matrix whose diagonal elements  $\lambda_i$  are the eigenvalues, and  $Z$  is a real orthogonal matrix whose columns  $z_i$  are the eigenvectors. This implies that  $z_i^T z_j = 0$  if  $i \neq j$ , and  $\|z_i\|_2 = 1$ .

Equation (2) can be rewritten

$$A = Z\Lambda Z^T. \quad (3)$$

This is known as the *eigen-decomposition* or *spectral factorization* of  $A$ .

Eigenvalues of a real symmetric matrix are well-conditioned, that is, they are not unduly sensitive to perturbations in the original matrix  $A$ . The sensitivity of an eigenvector depends on how small the gap is between its eigenvalue and any other eigenvalue: the smaller the gap, the more sensitive the eigenvector. More details on the accuracy of computed eigenvalues and eigenvectors are given in the routine documents, and in the F08 Chapter Introduction.

For dense or band matrices, the computation of eigenvalues and eigenvectors proceeds in the following stages:

1.  $A$  is reduced to a symmetric tridiagonal matrix  $T$  by an orthogonal similarity transformation:  $A = QTQ^T$ , where  $Q$  is orthogonal. (A *tridiagonal* matrix is zero except for the main diagonal and the first subdiagonal and superdiagonal on either side.)  $T$  has the same eigenvalues as  $A$  and is easier to handle.
2. Eigenvalues and eigenvectors of  $T$  are computed as required. If all eigenvalues (and optionally eigenvectors) are required, they are computed by the *QR* algorithm, which effectively factorizes  $T$  as  $T = SAS^T$ , where  $S$  is orthogonal, or by the divide-and-conquer method. If only selected eigenvalues are required, they are computed by bisection, and if selected eigenvectors are required, they are computed by inverse iteration. If  $s$  is an eigenvector of  $T$ , then  $Qs$  is an eigenvector of  $A$ .

All the above remarks also apply – with the obvious changes – to the case when  $A$  is a complex Hermitian matrix. The eigenvectors are complex, but the eigenvalues are all real, and so is the tridiagonal matrix  $T$ .

If  $A$  is large and sparse, the methods just described would be very wasteful in both storage and computing time, and therefore an alternative algorithm, known as *subspace iteration*, is provided (for real problems only) to find a (usually small) subset of the eigenvalues and their corresponding eigenvectors. Chapter F12 contains routines based on the Lanczos method for real symmetric large sparse eigenvalue problems, and these routines are usually more efficient than subspace iteration.

### 2.1.2 Standard nonsymmetric eigenvalue problems

A real nonsymmetric matrix  $A$  may have complex eigenvalues, occurring as complex conjugate pairs. If  $x$  is an eigenvector corresponding to a complex eigenvalue  $\lambda$ , then the complex conjugate vector  $\bar{x}$  is the eigenvector corresponding to the complex conjugate eigenvalue  $\bar{\lambda}$ . Note that the vector  $x$  defined in equation (1) is sometimes called a *right eigenvector*; a *left eigenvector*  $y$  is defined by

$$y^H A = \lambda y^H \quad \text{or} \quad A^T y = \bar{\lambda} y.$$

Routines in this chapter only compute right eigenvectors (the usual requirement), but routines in Chapter F08 can compute left or right eigenvectors or both.

The eigenvalue problem can be solved via the *Schur factorization* of  $A$ , defined as

$$A = ZTZ^T,$$

where  $Z$  is an orthogonal matrix and  $T$  is a real upper *quasi-triangular* matrix, with the same eigenvalues as  $A$ .  $T$  is called the *Schur form* of  $A$ . If all the eigenvalues of  $A$  are real, then  $T$  is upper triangular, and its diagonal elements are the eigenvalues of  $A$ . If  $A$  has complex conjugate pairs of eigenvalues, then  $T$  has 2 by 2 diagonal blocks, whose eigenvalues are the complex conjugate pairs of eigenvalues of  $A$ . (The structure of  $T$  is simpler if the matrices are complex – see below.)

For example, the following matrix is in quasi-triangular form

$$\begin{pmatrix} 1 & * & * & * \\ 0 & 2 & -1 & * \\ 0 & 1 & 2 & * \\ 0 & 0 & 0 & 3 \end{pmatrix}$$

and has eigenvalues 1,  $2 \pm i$ , and 3. (The elements indicated by ‘\*’ may take any values.)

The columns of  $Z$  are called the *Schur vectors*. For each  $k(1 \leq k \leq n)$ , the first  $k$  columns of  $Z$  form an orthonormal basis for the invariant subspace corresponding to the first  $k$  eigenvalues on the diagonal of  $T$ . (An *invariant subspace* (for  $A$ ) is a subspace  $S$  such that for any vector  $v$  in  $S$ ,  $Av$  is also in  $S$ .) Because this basis is orthonormal, it is preferable in many applications to compute Schur vectors rather than eigenvectors. It is possible to order the Schur factorization so that any desired set of  $k$  eigenvalues occupy the  $k$  leading positions on the diagonal of  $T$ , and routines for this purpose are provided in Chapter F08.

Note that if  $A$  is symmetric, the Schur vectors are the same as the eigenvectors, but if  $A$  is nonsymmetric, they are distinct, and the Schur vectors, being orthonormal, are often more satisfactory to work with in numerical computation.

Eigenvalues and eigenvectors of a nonsymmetric matrix may be ill-conditioned, that is, sensitive to perturbations in  $A$ . Chapter F08 contains routines which compute or estimate the condition numbers of eigenvalues and eigenvectors, and the F08 Chapter Introduction gives more details about the error analysis of nonsymmetric eigenproblems. The accuracy with which eigenvalues and eigenvectors can be obtained is often improved by *balancing* a matrix. This is discussed further in Section 3.4.

Computation of eigenvalues, eigenvectors or the Schur factorization proceeds in the following stages:

1.  $A$  is reduced to an upper Hessenberg matrix  $H$  by an orthogonal similarity transformation:  $A = QHQ^T$ , where  $Q$  is orthogonal. (An *upper Hessenberg* matrix is zero below the first subdiagonal.)  $H$  has the same eigenvalues as  $A$ , and is easier to handle.
2. The upper Hessenberg matrix  $H$  is reduced to Schur form  $T$  by the  $QR$  algorithm, giving the Schur factorization  $H = STS^T$ . The eigenvalues of  $A$  are obtained from the diagonal blocks of  $T$ . The matrix  $Z$  of Schur vectors (if required) is computed as  $Z = QS$ .
3. After the eigenvalues have been found, eigenvectors may be computed, if required, in two different ways. Eigenvectors of  $H$  can be computed by inverse iteration, and then pre-multiplied by  $Q$  to give eigenvectors of  $A$ ; this approach is usually preferred if only a few eigenvectors are required. Alternatively, eigenvectors of  $T$  can be computed by back-substitution, and pre-multiplied by  $Z$  to give eigenvectors of  $A$ .

All the above remarks also apply – with the obvious changes – to the case when  $A$  is a complex matrix. The eigenvalues are in general complex, so there is no need for special treatment of complex conjugate pairs, and the Schur form  $T$  is simply a complex upper triangular matrix.

As for the symmetric eigenvalue problem, if  $A$  is large and sparse then it is generally preferable to use an alternative method. Chapter F12 provides routines based on Arnoldi's method for both real and complex matrices, intended to find a subset of the eigenvalues and vectors.

## 2.2 The Singular Value Decomposition

The *singular value decomposition* (SVD) of a real  $m$  by  $n$  matrix  $A$  is given by

$$A = U\Sigma V^T,$$

where  $U$  and  $V$  are orthogonal and  $\Sigma$  is an  $m$  by  $n$  diagonal matrix with real diagonal elements,  $\sigma_i$ , such that

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{\min(m,n)} \geq 0.$$

The  $\sigma_i$  are the *singular values* of  $A$  and the first  $\min(m, n)$  columns of  $U$  and  $V$  are, respectively, the *left* and *right singular vectors* of  $A$ . The singular values and singular vectors satisfy

$$Av_i = \sigma_i u_i \quad \text{and} \quad A^T u_i = \sigma_i v_i$$

where  $u_i$  and  $v_i$  are the  $i$ th columns of  $U$  and  $V$  respectively.

The singular value decomposition of  $A$  is closely related to the eigen-decompositions of the symmetric matrices  $A^T A$  or  $AA^T$ , because:

$$A^T Av_i = \sigma_i^2 v_i \quad \text{and} \quad AA^T u_i = \sigma_i^2 u_i.$$

However, these relationships are not recommended as a means of computing singular values or vectors unless  $A$  is sparse and routines from Chapter F12 are to be used.

If  $U_k, V_k$  denote the leading  $k$  columns of  $U$  and  $V$  respectively, and if  $\Sigma_k$  denotes the leading principal submatrix of  $\Sigma$ , then

$$A_k \equiv U_k \Sigma_k V_k^T$$

is the best rank- $k$  approximation to  $A$  in both the 2-norm and the Frobenius norm.

Singular values are well-conditioned; that is, they are not unduly sensitive to perturbations in  $A$ . The sensitivity of a singular vector depends on how small the gap is between its singular value and any other singular value: the smaller the gap, the more sensitive the singular vector. More details on the accuracy of computed singular values and vectors are given in the routine documents and in the F08 Chapter Introduction.

The singular value decomposition is useful for the numerical determination of the rank of a matrix, and for solving linear least squares problems, especially when they are rank-deficient (or nearly so). See Chapter F04.

Computation of singular values and vectors proceeds in the following stages:

1.  $A$  is reduced to an upper bidiagonal matrix  $B$  by an orthogonal transformation  $A = U_1 B V_1^T$ , where  $U_1$  and  $V_1$  are orthogonal. (An *upper bidiagonal* matrix is zero except for the main diagonal and the first superdiagonal.)  $B$  has the same singular values as  $A$ , and is easier to handle.
2. The SVD of the bidiagonal matrix  $B$  is computed as  $B = U_2 \Sigma V_2^T$ , where  $U_2$  and  $V_2$  are orthogonal and  $\Sigma$  is diagonal as described above. Then in the SVD of  $A$ ,  $U = U_1 U_2$  and  $V = V_1 V_2$ .

All the above remarks also apply – with the obvious changes – to the case when  $A$  is a complex matrix. The singular vectors are complex, but the singular values are real and non-negative, and the bidiagonal matrix  $B$  is also real.

By formulating the problems appropriately, real large sparse singular value problems may be solved using the symmetric eigenvalue routines in Chapter F12.

## 2.3 Generalized Eigenvalue Problems

Let  $A$  and  $B$  be square matrices of order  $n$ . The *generalized eigenvalue problem* is to find eigenvalues,  $\lambda$ , and corresponding eigenvectors,  $x \neq 0$ , such that

$$Ax = \lambda Bx. \quad (4)$$

For given  $A$  and  $B$ , the set of all matrices of the form  $A - \lambda B$  is called a *pencil*, and  $\lambda$  and  $x$  are said to be an eigenvalue and eigenvector of the pencil  $A - \lambda B$ .

When  $B$  is nonsingular, equation (4) is mathematically equivalent to  $(B^{-1}A)x = \lambda x$ , and when  $A$  is nonsingular, it is equivalent to  $(A^{-1}B)x = (1/\lambda)x$ . Thus, in theory, if one of the matrices  $A$  or  $B$  is known to be nonsingular, the problem could be reduced to a standard eigenvalue problem.

However, for this reduction to be satisfactory from the point of view of numerical stability, it is necessary not only that  $B$  (or  $A$ ) should be nonsingular, but that it should be well-conditioned with respect to inversion. The nearer  $B$  is to singularity, the more unsatisfactory  $B^{-1}A$  will be as a vehicle for determining the required eigenvalues. Well-determined eigenvalues of the original problem (4) may be poorly determined even by the correctly rounded version of  $B^{-1}A$ .

We consider first a special class of problems in which  $B$  is known to be nonsingular, and then return to the general case in the following sub-section.

### 2.3.1 Generalized symmetric-definite eigenvalue problems

If  $A$  and  $B$  are *symmetric* and  $B$  is *positive definite*, then the generalized eigenvalue problem has desirable properties similar to those of the standard symmetric eigenvalue problem. The eigenvalues are all real, and the eigenvectors, while not orthogonal in the usual sense, satisfy the relations  $z_i^T B z_j = 0$  for  $i \neq j$  and can be normalized so that  $z_i^T B z_i = 1$ .

Note that it is not enough for  $A$  and  $B$  to be symmetric;  $B$  must also be positive definite, which implies nonsingularity. Eigenproblems with these properties are referred to as *symmetric-definite* problems.

If  $A$  is the diagonal matrix whose diagonal elements are the eigenvalues, and  $Z$  is the matrix whose columns are the eigenvectors, then

$$Z^T A Z = A \quad \text{and} \quad Z^T B Z = I.$$

To compute eigenvalues and eigenvectors, the problem can be reduced to a standard symmetric eigenvalue problem, using the Cholesky factorization of  $B$  as  $LL^T$  or  $U^T U$  (see Chapter F07). Note, however, that this reduction does implicitly involve the inversion of  $B$ , and hence this approach should *not* be used if  $B$  is ill-conditioned with respect to inversion.

For example, with  $B = LL^T$ , we have

$$Az = \lambda Bz \Leftrightarrow (L^{-1} A L^{-T})(L^T z) = \lambda(L^T z).$$

Hence the eigenvalues of  $Az = \lambda Bz$  are those of  $Cy = \lambda y$ , where  $C$  is the symmetric matrix  $C = L^{-1} A L^{-T}$  and  $y = L^T z$ . The standard symmetric eigenproblem  $Cy = \lambda y$  may be solved by the methods described in Section 2.1.1. The eigenvectors  $z$  of the original problem may be recovered by computing  $z = L^{-T} y$ .

Most of the routines which solve this class of problems can also solve the closely related problems

$$ABx = \lambda x \quad \text{or} \quad BAx = \lambda x$$

where again  $A$  and  $B$  are symmetric and  $B$  is positive definite. See the routine documents for details.

All the above remarks also apply – with the obvious changes – to the case when  $A$  and  $B$  are complex Hermitian matrices. Such problems are called *Hermitian-definite*. The eigenvectors are complex, but the eigenvalues are all real.

If  $A$  and  $B$  are large and sparse, reduction to an equivalent standard eigenproblem as described above would almost certainly result in a large dense matrix  $C$ , and hence would be very wasteful in both storage and computing time. The methods of subspace iteration and Lanczos type methods, mentioned in Section 2.1.1, can also be used for large sparse generalized symmetric-definite problems.

### 2.3.2 Generalized nonsymmetric eigenvalue problems

Any generalized eigenproblem which is not symmetric-definite with well-conditioned  $B$  must be handled as if it were a general nonsymmetric problem.

If  $B$  is singular, the problem has infinite eigenvalues. These are not a problem; they are equivalent to zero eigenvalues of the problem  $Bx = \mu Ax$ . Computationally they appear as very large values.

If  $A$  and  $B$  are both singular and have a common null space, then  $A - \lambda B$  is singular for all  $\lambda$ ; in other words, any value  $\lambda$  can be regarded as an eigenvalue. Pencils with this property are called *singular*.

As with standard nonsymmetric problems, a real problem may have complex eigenvalues, occurring as complex conjugate pairs.

The generalized eigenvalue problem can be solved via the *generalized Schur factorization* of  $A$  and  $B$ :

$$A = Q U Z^T, \quad B = Q V Z^T$$

where  $Q$  and  $Z$  are orthogonal,  $V$  is upper triangular, and  $U$  is upper quasi-triangular (defined just as in Section 2.1.2).

If all the eigenvalues are real, then  $U$  is upper triangular; the eigenvalues are given by  $\lambda_i = u_{ii}/v_{ii}$ . If there are complex conjugate pairs of eigenvalues, then  $U$  has 2 by 2 diagonal blocks.

Eigenvalues and eigenvectors of a generalized nonsymmetric problem may be ill-conditioned; that is, sensitive to perturbations in  $A$  or  $B$ .

Particular care must be taken if, for some  $i$ ,  $u_{ii} = v_{ii} = 0$ , or in practical terms if  $u_{ii}$  and  $v_{ii}$  are both small; this means that the pencil is singular, or approximately so. Not only is the particular value  $\lambda_i$

undetermined, but also **no reliance** can be placed on **any** of the computed eigenvalues. See also the routine documents.

Computation of eigenvalues and eigenvectors proceeds in the following stages.

1. The pencil  $A - \lambda B$  is reduced by an orthogonal transformation to a pencil  $H - \lambda K$  in which  $H$  is upper Hessenberg and  $K$  is upper triangular:  $A = Q_1 H Z_1^T$  and  $B = Q_1 K Z_1^T$ . The pencil  $H - \lambda K$  has the same eigenvalues as  $A - \lambda B$ , and is easier to handle.
2. The upper Hessenberg matrix  $H$  is reduced to upper quasi-triangular form, while  $K$  is maintained in upper triangular form, using the  $QZ$  algorithm. This gives the generalized Schur factorization:  $H = Q_2 U Z_2$  and  $K = Q_2 V Z_2$ .
3. Eigenvectors of the pencil  $U - \lambda V$  are computed (if required) by back-substitution, and pre-multiplied by  $Z_1 Z_2$  to give eigenvectors of  $A$ .

All the above remarks also apply – with the obvious changes – to the case when  $A$  and  $B$  are complex matrices. The eigenvalues are in general complex, so there is no need for special treatment of complex conjugate pairs, and the matrix  $U$  in the generalized Schur factorization is simply a complex upper triangular matrix.

As for the generalized symmetric-definite eigenvalue problem, if  $A$  and  $B$  are large and sparse then it is generally preferable to use an alternative method. Chapter F12 provides routines based on Arnoldi's method for both real and complex matrices, intended to find a subset of the eigenvalues and vectors.

## 2.4 Quadratic eigenvalue problems

Let  $A$ ,  $B$  and  $C$  be square matrices of order  $n$ . The *quadratic eigenvalue problem* (QEP) is to find eigenvalues,  $\lambda$ , and corresponding eigenvectors,  $x \neq 0$ , such that

$$(\lambda^2 A + \lambda B + C)x = 0.$$

More specifically,  $x$  is a right eigenvector and a left eigenvector,  $y$ , is such that

$$y^H(\lambda^2 A + \lambda B + C) = 0,$$

where  $y^H$  is the conjugate transpose of  $y$  (transpose when  $y$  is real).

In general the QEP has  $2n$  eigenvalues and corresponding eigenvectors.

QEPs are generally solved by linearizing the problem to produce a  $2n$  by  $2n$  generalized eigenvalue problem. For example,

$$C_1(\lambda) = \begin{pmatrix} B & C \\ -I & 0 \end{pmatrix} - \lambda \begin{pmatrix} -A & 0 \\ 0 & -I \end{pmatrix},$$

which is called the first companion form and has the same  $2n$  eigenvalues as the QEP.

If

$$\det(\lambda^2 A + \lambda B + C) \neq 0,$$

then the QEP is said to be *regular*, or *non-singular*. For a regular QEP, when  $C$  is singular the QEP has one or more zero eigenvalues and when  $A$  is singular the QEP has one or more infinite eigenvalues. As with the generalized problem particular care must be taken when the problem is singular (see Section 2.3.2).

As with generalized nonsymmetric problems, a real QEP may have complex eigenvalues, occurring as complex conjugate pairs.

For further information on QEPs, see the survey article by Tisseur and Meerbergen (2001), and for details on solving the dense QEP see Hammarling *et al.* (2013).

## 3 Recommendations on Choice and Use of Available Routines

### 3.1 Black Box Routines and General Purpose Routines

Routines in the NAG Library for solving eigenvalue problems fall into two categories.

1. **Black Box Routines:** these are designed to solve a standard type of problem in a single call – for example, to compute all the eigenvalues and eigenvectors of a real symmetric matrix. You are recommended to use a black box routine if there is one to meet your needs; refer to the decision tree in Section 4.1 or the index in Section 5.
2. **General Purpose Routines:** these perform the computational subtasks which make up the separate stages of the overall task, as described in Section 2 – for example, reducing a real symmetric matrix to tridiagonal form. General purpose routines are to be found, for historical reasons, some in this chapter, a few in Chapter F01, but most in Chapter F08. If there is no black box routine that meets your needs, you will need to use one or more general purpose routines.

The decision trees in Section 4.2 list the combinations of general purpose routines which are needed to solve many common types of problem.

Sometimes a combination of a black box routine and one or more general purpose routines will be the most convenient way to solve your problem: the black box routine can be used to compute most of the results, and a general purpose routine can be used to perform a subsidiary computation, such as computing condition numbers of eigenvalues and eigenvectors.

### 3.2 Computing Selected Eigenvalues and Eigenvectors

The decision trees and the routine documents make a distinction between routines which compute *all* eigenvalues or eigenvectors, and routines which compute *selected* eigenvalues or eigenvectors; the two classes of routine use different algorithms.

It is difficult to give clear guidance on which of these two classes of routine to use in a particular case, especially with regard to computing eigenvectors. If you only wish to compute a very few eigenvectors, then a routine for selected eigenvectors will be more economical, but if you want to compute a substantial subset (an old rule of thumb suggested more than 25%), then it may be more economical to compute all of them. Conversely, if you wish to compute all the eigenvectors of a sufficiently large symmetric tridiagonal matrix, the routine for selected eigenvectors may be faster.

The choice depends on the properties of the matrix and on the computing environment; if it is critical, you should perform your own timing tests.

For dense nonsymmetric eigenproblems, there are no algorithms provided for computing selected eigenvalues; it is always necessary to compute all the eigenvalues, but you can then select specific eigenvectors for computation by inverse iteration.

### 3.3 Storage Schemes for Symmetric Matrices

Routines which handle symmetric matrices are usually designed to use either the upper or lower triangle of the matrix; it is not necessary to store the whole matrix. If either the upper or lower triangle is stored conventionally in the upper or lower triangle of a two-dimensional array, the remaining elements of the array can be used to store other useful data. However, that is not always convenient, and if it is important to economize on storage, the upper or lower triangle can be stored in a one-dimensional array of length  $n(n+1)/2$ ; in other words, the storage is almost halved. This storage format is referred to as *packed storage*.

Routines designed for packed storage are usually less efficient, especially on high-performance computers, so there is a trade-off between storage and efficiency.

A *band* matrix is one whose nonzero elements are confined to a relatively small number of subdiagonals or superdiagonals on either side of the main diagonal. Algorithms can take advantage of bandedness to reduce the amount of work and storage required.

Routines which take advantage of packed storage or bandedness are provided for both standard symmetric eigenproblems and generalized symmetric-definite eigenproblems.



### 3.4 Balancing for Nonsymmetric Eigenproblems

There are two preprocessing steps which one may perform on a nonsymmetric matrix  $A$  in order to make its eigenproblem easier. Together they are referred to as *balancing*.

1. **Permutation:** this involves reordering the rows and columns to make  $A$  more nearly upper triangular (and thus closer to Schur form):  $A' = PAP^T$ , where  $P$  is a permutation matrix. If  $A$  has a significant number of zero elements, this preliminary permutation can reduce the amount of work required, and also improve the accuracy of the computed eigenvalues. In the extreme case, if  $A$  is permutable to upper triangular form, then no floating-point operations are needed to reduce it to Schur form.
2. **Scaling:** a diagonal matrix  $D$  is used to make the rows and columns of  $A'$  more nearly equal in norm:  $A'' = DA'D^{-1}$ . Scaling can make the matrix norm smaller with respect to the eigenvalues, and so possibly reduce the inaccuracy contributed by roundoff (see Chapter II/11 of Wilkinson and Reinsch (1971)).

Routines are provided in Chapter F08 for performing either or both of these preprocessing steps, and also for transforming computed eigenvectors or Schur vectors back to those of the original matrix.

The black box routines in this chapter which compute eigenvectors perform both forms of balancing.

### 3.5 Non-uniqueness of Eigenvectors and Singular Vectors

Eigenvectors, as defined by equations (1) or (4), are *not uniquely defined*. If  $x$  is an eigenvector, then so is  $kx$  where  $k$  is any nonzero scalar. Eigenvectors computed by different algorithms, or on different computers, may appear to disagree completely, though in fact they differ only by a scalar factor (which may be complex). These differences should not be significant in any application in which the eigenvectors will be used, but they can arouse uncertainty about the correctness of computed results.

Even if eigenvectors  $x$  are normalized so that  $\|x\|_2 = 1$ , this is not sufficient to fix them uniquely, since they can still be multiplied by a scalar factor  $k$  such that  $|k| = 1$ . To counteract this inconvenience, most of the routines in this chapter, and in Chapter F08, normalize eigenvectors (and Schur vectors) so that  $\|x\|_2 = 1$  and the component of  $x$  with largest absolute value is real and positive. (There is still a possible indeterminacy if there are two components of equal largest absolute value – or in practice if they are very close – but this is rare.)

In symmetric problems the computed eigenvalues are sorted into ascending order, but in nonsymmetric problems the order in which the computed eigenvalues are returned is dependent on the detailed working of the algorithm and may be sensitive to rounding errors. The Schur form and Schur vectors depend on the ordering of the eigenvalues and this is another possible cause of non-uniqueness when they are computed. However, it must be stressed again that variations in the results from this cause should not be significant. (Routines in Chapter F08 can be used to transform the Schur form and Schur vectors so that the eigenvalues appear in any given order if this is important.)

In singular value problems, the left and right singular vectors  $u$  and  $v$  which correspond to a singular value  $\sigma$  cannot be normalized independently: if  $u$  is multiplied by a factor  $k$  such that  $|k| = 1$ , then  $v$  must also be multiplied by  $k$ .

Non-uniqueness also occurs among eigenvectors which correspond to a multiple eigenvalue, or among singular vectors which correspond to a multiple singular value. In practice, this is more likely to be apparent as the extreme sensitivity of eigenvectors which correspond to a cluster of close eigenvalues (or of singular vectors which correspond to a cluster of close singular values).

## 4 Decision Trees

### 4.1 Black Box Routines

The decision tree for this section is divided into three sub-trees.


Tree 1 Eigenvalues and Eigenvectors of Real Matrices

Tree 2 Eigenvalues and Eigenvectors of Complex Matrices


Tree 3 Singular Values and Singular Vectors

**Note:** for the Chapter F08 routines there is generally a choice of simple and comprehensive routine. The comprehensive routines return additional information such as condition and/or error estimates.


**Tree 1: Eigenvalues and Eigenvectors of Real Matrices**



**Tree 2: Eigenvalues and Eigenvectors of Complex Matrices**



**Tree 3: Singular Values and Singular Vectors**



**4.2 General Purpose Routines (Eigenvalues and Eigenvectors)**

Routines for large sparse eigenvalue problems are to be found in Chapter F12, see the F12 Chapter Introduction.

The decision tree for this section addressing dense problems, is divided into eight sub-trees:

- Tree 1 Real Symmetric Eigenvalue Problems in the F08 Chapter Introduction
- Tree 2 Real Generalized Symmetric-definite Eigenvalue Problems in the F08 Chapter Introduction
- Tree 3 Real Nonsymmetric Eigenvalue Problems in the F08 Chapter Introduction
- Tree 4 Real Generalized Nonsymmetric Eigenvalue Problems in the F08 Chapter Introduction
- Tree 5 Complex Hermitian Eigenvalue Problems in the F08 Chapter Introduction

Tree 6 Complex Generalized Hermitian-definite Eigenvalue Problems in the F08 Chapter Introduction

Tree 7 Complex non-Hermitian Eigenvalue Problems in the F08 Chapter Introduction

Tree 8 Complex Generalized non-Hermitian Eigenvalue Problems in the F08 Chapter Introduction

As it is very unlikely that one of the routines in this section will be called on its own, the other routines required to solve a given problem are listed in the order in which they should be called.

### 4.3 General Purpose Routines (Singular Value Decomposition)

See Section 4.2 in the F08 Chapter Introduction. For real sparse matrices where only selected singular values are required (possibly with their singular vectors), routines from Chapter F12 may be applied to the symmetric matrix  $A^T A$ ; see Section 10 in F12FBF.

## 5 Functionality Index

Black Box routines,

complex eigenproblem, selected eigenvalues and eigenvectors .....	F02GCF
complex quadratic eigenproblem, all eigenvalues and optionally eigenvectors, backward, errors and eigenvalue condition numbers .....	F02JQF
complex upper triangular matrix, singular values and, optionally, left and/or right singular vectors .....	F02XUF
generalized real sparse symmetric-definite eigenproblem, selected eigenvalues and eigenvectors .....	F02FJF
real eigenproblem, selected eigenvalues and eigenvectors .....	F02ECF
real quadratic eigenproblem, all eigenvalues and optionally eigenvectors, backward, errors and eigenvalue condition numbers .....	F02JCF
real sparse eigenproblem, selected eigenvalues and eigenvectors .....	F02EKF
real sparse symmetric matrix, driver, selected eigenvalues and eigenvectors .....	F02FKF
selected eigenvalues and eigenvectors .....	F02FJF
real upper triangular matrix, singular values and, optionally, left and/or right singular vectors .....	F02WUF
General Purpose routines (see also Chapter F12), real $m$ by $n$ matrix, leading terms SVD .....	F02WGF

## 6 Auxiliary Routines Associated with Library Routine Parameters

F02EKY	nagf_eigen_arnoldi_option See the description of the parameter OPTION in F02EKF.
F02EKZ	nagf_eigen_arnoldi_monit_gen See the description of the parameter MONIT in F02EKF.
F02FJZ	nagf_eigen_monit See the description of the parameter MONIT in F02FJF.
F02FKZ	nagf_eigen_arnoldi_monit_symm See the description of the parameter MONIT in F02FKF.

## 7 Routines Withdrawn or Scheduled for Withdrawal

The following lists all those routines that have been withdrawn since Mark 18 of the Library or are scheduled for withdrawal at one of the next two marks.

<b>Withdrawn Routine</b>	<b>Mark of Withdrawal</b>	<b>Replacement Routine(s)</b>
F02AAF	18	F08FAF (DSYEV)
F02ABF	18	F08FAF (DSYEV)
F02ADF	18	F08SAF (DSYGV)
F02AEF	18	F08SAF (DSYGV)
F02AFF	18	F08NAF (DGEEV)
F02AGF	18	F08NAF (DGEEV)
F02AJF	18	F08NNF (ZGEEV)
F02AKF	18	F08NNF (ZGEEV)
F02AMF	18	F08JEF (DSTEQR)
F02ANF	18	F08PSF (ZHSEQR)
F02APF	18	F08PEF (DHSEQR)
F02AQF	18	F08PEF (DHSEQR) and F08QKF (DTREVC)
F02ARF	18	F08PSF (ZHSEQR) and F08QXF (ZTREVC)
F02AVF	18	F08JFF (DSTERF)
F02AWF	18	F08FNF (ZHEEV)
F02AXF	18	F08FNF (ZHEEV)
F02AYF	18	F08JSF (ZSTEQR)
F02BBF	19	F08FBF (DSYEVX)
F02BCF	19	F02ECF
F02BDF	19	F02GCF
F02BEF	18	F08JFF (DSTEBZ) and F08JKF (DSTEIN)
F02BFF	18	F08JFF (DSTEBZ)
F02BJF	23	F08WAF (DGGEV)
F02BKF	18	F08PKF (DHSEIN)
F02BLF	18	F08PXF (ZHSEIN)
F02EAF	23	F08PAF (DGEEV)
F02EBF	23	F08NAF (DGEEV)
F02FAF	23	F08FAF (DSYEV)
F02FCF	23	F08FBF (DSYEVX)
F02FDF	23	F08SAF (DSYGV)
F02FHF	23	F08UAF (DSBGV)
F02GAF	23	F08PNF (ZGEES)
F02GBF	23	F08NNF (ZGEEV)
F02GJF	23	F08WNF (ZGGEV)
F02HAF	23	F08FNF (ZHEEV)
F02HCF	23	F08FPF (ZHEEVX)
F02HDF	23	F08SNF (ZHEGV)
F02SDF	27	F12AGF and F12FGF
F02SWF	18	F08KEF (DGEBRD)
F02SXF	18	F08KFF (DORGBR) and F08KGF (DORMBR)
F02SYF	18	F08MEF (DBDSQR)
F02UWF	18	F08KSF (ZGEBRD), F06TFF, F08KTF (ZUNGBR) and F08KUF (ZUNMBR)
F02UXF	18	F08KTF (ZUNGBR) or F08KUF (ZUNMBR)
F02UYF	18	F08MSF (ZBDSQR)
F02WDF	27	F02WUF and F08AEF (DGEQRF)
F02WEF	23	F08KBF (DGESVD)
F02XEF	23	F08KPF (ZGESVD)

## 8 References

Anderson E, Bai Z, Bischof C, Blackford S, Demmel J, Dongarra J J, Du Croz J J, Greenbaum A, Hammarling S, McKenney A and Sorensen D (1999) *LAPACK Users' Guide* (3rd Edition) SIAM, Philadelphia

Golub G H and Van Loan C F (1996) *Matrix Computations* (3rd Edition) Johns Hopkins University Press, Baltimore

Hammarling S, Munro C J and Tisseur F (2013) An algorithm for the complete solution of quadratic eigenvalue problems. *ACM Trans. Math. Software.* **39(3):18:1–18:119** (<http://eprints.ma.man.ac.uk/1815/>)

Parlett B N (1998) *The Symmetric Eigenvalue Problem* SIAM, Philadelphia

Tisseur F and Meerbergen K (2001) The quadratic eigenvalue problem *SIAM Review* **43:235–286**

Wilkinson J H and Reinsch C (1971) *Handbook for Automatic Computation II, Linear Algebra* Springer-Verlag

---